



Cardiology and Angiology: An International Journal

7(3): 1-8, 2018; Article no.CA.40399
ISSN: 2347-520X, NLM ID: 101658392

Cardiovascular Disease Risk Factors among Older Prisoners in the Ashanti Region of Ghana

**Nana Ama Frimpomaa Agyapong^{1*}, Reginald Adjetey Annan¹
and Charles Apprey¹**

¹*Department of Biochemistry and Biotechnology, College of Science, Kwame Nkrumah University of Science and Technology, Ghana.*

Authors' contributions

This work was carried out in collaboration between all authors. Authors NAFA and RAA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors RAA and CA managed the analyses of the study. Authors CA and NAFA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CA/2018/40399

Editor(s):

(1) Francesco Pelliccia, Professor, Department of Heart and Great Vessels, University La Sapienza, Rome, Italy.

Reviewers:

(1) Anuj Nautiyal, DIT University, India.

(2) Anonymous, Usmanu Danfodiyo University, Nigeria.

Complete Peer review History: <http://www.sciencedomain.org/review-history/23830>

Original Research Article

Received 17th January 2018
Accepted 23rd March 2018
Published 27th March 2018

ABSTRACT

Prisons are environments that have been characterised by high rates of communicable diseases until recently when the prevalence of cardiovascular diseases have been assessed, and high prevalence found. This study, involving 160 inmates assessed the prevalence of cardiovascular risk factors among prisoners in the Ashanti Region of Ghana. BMI, waist circumference and blood pressure of all participants were checked. Additionally, lifestyle and dietary factors such as exercise and fruit intake as well as medical history of inmates were assessed. Fasting blood samples were taken and analysed for lipid profile and FBG. The mean systolic blood pressure of study participants was 141.1±23.2 mmHg, diastolic 88.9±15 mmHg, BMI 22.8±4.1 kg/m², waist circumference 81±10.3 cm, FBG 4.3±0.9 mmol/L, HDL 1.4±0.4 mmol/L and triglycerides 1.1±0.6 mmol/L. Prevalence of hypertension and dyslipidaemia was 57.5% each. Prevalence of metabolic syndrome was 8.1%, and 21.9% had two cardiovascular risk factors. In conclusion prevalence of dyslipidaemia and elevated blood pressure were high among inmates and cardiovascular risk factors were higher among female

*Corresponding author: E-mail: nagyapong29@gmail.com;

inmates compared to males. Most inmates were sedentary and occasionally consumed fruits. Interventions of appropriate dietary provision and exercise schedule should begin within Ghanaian prisons to curb this menace.

Keywords: Cardiovascular diseases; metabolic syndrome; inmates; prisons; risk factors.

1. INTRODUCTION

Cardiovascular diseases are a significant public health concern, and their occurrence is firmly attributed to modifiable risk factors [1]. These modifiable risk factors include elevated blood pressure, diabetes or glucose intolerance, dyslipidaemia and central obesity [2]. According to the National Cholesterol Education Programme, Adult Treatment Panel III, the presence of at least any three of these factors within an individual satisfies the diagnostic criteria for metabolic syndrome and it is a significant predictor of future mortality and morbidity [3].

Prisons are stressful environments with meal provisions that predisposes to cardiovascular diseases [4,5]. In a study conducted in Istanbul, it was documented that cardiovascular diseases are the most common cause of natural death among prisoners [6]. Additionally, in the United States, prisoners had a higher risk of cardiovascular diseases compared to the general population even with adjustments for important socio-demographic factors [7]. Inappropriate diet, smoking, high alcohol consumption as well as physical inactivity have been outlined as the primary causes of metabolic risk markers within prisons [8].

The 2012 annual report on inmate mortality by the Ghana Prisons Service indicates that cardiovascular disease is the major cause of mortality among Ghanaian inmates.

Older inmates aged forty (40) and above are of increasing concern with regards to cardiovascular risk and health status compared to younger ones [9]. This may be because increasing age is an independent risk factor for cardiovascular disease coupled with the stressful prison environment and inappropriate diet [10,11]. High prevalence of cardiovascular diseases among inmates adds to the cost of their basic care which is already a burden for developing countries like Ghana but early assessment of these risk factors can enable timely interventions to be implemented [12-14].

The primary aim of this study was to assess the prevalence of cardiovascular risk factors among older prisoners in the Ashanti Region of Ghana.

2. METHODS

2.1 Study Population

A sample size of one hundred and sixty was used for the study to make findings more applicable to the inmate population. The Cochran's formula was used to estimate the minimal sample size (Cochran, 1977). Below is the calculation;

$$\text{Sample size} = Z_{1-\alpha/2}^2 p(1-p)/d^2$$

$$Z_{1-\alpha/2}^2 = 1.96 \text{ for confidence interval of } 95\%$$

p = Expected proportion in a population based on previous reports is 0.11

d = absolute error of precision which taken as 0.05 for this study.

$$\text{Sample size} = 1.96^2 \times 0.11(1-0.11)/0.05^2$$

$$\text{Sample size} = 3.8416 \times 0.11(0.89)/0.0025$$

$$\text{Sample size} = 151$$

The 160 inmates were from the Kumasi central (131), Kumasi female (10) and Manhyia local prisons (19) all in the Ashanti Region of Ghana. Prison officers helped in the selection of inmates who met the inclusion criteria. In all, the study was explained to 169 inmates who qualified to partake in the study but only 160 consented to participate.

2.2 Inclusion and Exclusion Criteria

Inmates aged forty and above who had been within prison for more than three months formed the inclusion criteria. Inmates who were less than forty years and those who were ill were excluded.

2.3 Data Collection

A detailed questionnaire was used to collect information on socio-demographic characteristics of inmates, medical history and voluntary

exercise. About 5 ml of fasting blood sample of each participant was taken and analysed for lipid profile and FBG. About 2 mls of the sample taken was dispensed into a fluoride tube for glucose analysis and the remaining sample into an activator gel tube for the analysis of lipids. The blood samples were kept in an ice chest containing ice packs and transported to the clinical analysis laboratory of KNUST for the biochemical analysis. The samples for lipid profile analysis were centrifuged for ten minutes at a speed of 4000 rotation per minute (r.p.m) using the eppendorf centrifuge 5804 to obtain the serum for the analysis. Lipid profile and FBG were analysed using the Randox rx monza semi-automated spectrophotometer. FBG (mmol/L) was used to classify inmates as hypoglycaemic (>3.5), normoglycaemic (>5.6), pre-diabetic (5.6-7.0) and diabetic (>7.0) (ADA, 2010). Serum low density lipoprotein cholesterol (LDL-c) of (≥ 4.12 mmol/L) and total cholesterol (TC) of (≥ 5.18) were classified as high [15]. Weight and height were also measured and used to calculate BMI (weight/height in metres²). Waist circumference of all inmates was taken with a plastic tape and systolic and diastolic blood pressure of study participants was taken twice using a digital sphygmomanometer. The latter reading was used for the analysis. This was used to classify participants as normotensive (< 120/80 mm Hg), pre hypertensive (120-139 mm Hg systolic and or diastolic 80-89 mm Hg) and hypertensive (> 140/90 mm Hg).

Metabolic syndrome was characterized by the components defined by the NCEP ATP III. These components include central obesity (waist circumference ≥ 102 cm in men and ≥ 88 cm in women), elevated blood pressure (≥ 130 mmHg systolic and or ≥ 85 mmHg diastolic), impaired fasting plasma glucose (≥ 5.6 mmol/L), decreased high density lipoproteins cholesterol (HDL-c) (< 1.03 mmol/L) and elevated triglycerides (≥ 1.7 mmol/L). BMI was used to categorize inmates as underweight (>18.5 kg/m²), normal (18.5-24.9 kg/m²), overweight (25-29.9 kg/m²) and obese (>30 kg/m²) [16].

2.4 Ethical Consideration

Ethical approval for the study was granted by the Committee on Human Research Publication and Ethics (CHRPE), School of Medical Sciences, KNUST, Kumasi; ref CHRPE/AP/407/16. Approval was also sought and granted by the Ghana Prisons Service headquarters before data collection at the prison facilities. Inmates who did not consent to the study were excluded and

those that consented signed or thumb-printed a consent form.

2.5 Data Analysis

IBM SPSS version 20 was used for the analysis of data. Student T- test was used to compare the means of parameters of the study and chi-square was used to compare categorical data. A p-value of <0.05 was set as statistically significant.

3. RESULTS

A total of one hundred and sixty (160) prisoners took part in the study of which 10 (6.3%) were females and the rest were males (93.8%). With respect to age distribution, 40.6% of the inmates were within the age range of 40-45, 22.5% within 46-50, 15.6% within 51-55, 6.9% within 56-60 and 14.4% were more than 60 years old. Table 1 displays the socio-demographic and incarceration characteristics of inmates.

Table 2 shows the means of metabolic parameters. The mean systolic and diastolic blood pressure of all study participants was 141.1 \pm 23.2 mm Hg and 88.9 \pm 14 mm Hg respectively. Mean BMI and waist circumference were 22.8 \pm 4.1 kg/m² and 81.1 \pm 10.3 cm respectively for all participants. Differences in BMI, diastolic blood pressure, FBG and triglycerides were statistically significant between males and females with females recording higher means for all with the exception of triglycerides.

Table 3 shows the metabolic characteristics of study participants. The overall prevalence of hypertension was 57.5% among all inmates followed by pre-hypertension (28.8%). Normotensive subjects constituted 13.8% of the total study population. Data on lipid profile was missing for one person. In total, those with dyslipidaemia constituted 57.5% of the study population. Prevalence of pre-diabetes, hypertension, overweight, obesity and metabolic syndrome was significantly higher in females compared to males.

Table 4 shows the physical activity levels of inmates. Most inmates were sedentary (55%) and those who exercised only did so occasionally (13.8%).

Table 5 shows the past medical history of inmates. Inmates with known hypertension constituted 11.9%. Out of this percentage, 10.6% were on medication but none was on special diet.

Table 1. Socio-demographics and incarceration characteristics of participants

Variable	Total n (%)
Gender	
Male	150 (93.8)
Female	10 (6.3)
Age	
40-45 years	65 (40.6)
46-50 years	36 (22.5)
51-55 years	25 (15.6)
56-60 years	11 (6.9)
>60 years	23 (14.4)
Education	
None	30 (18.8)
JHS*	85 (53.1)
SHS*	38 (23.8)
Tertiary	7 (4.4)
Marital status	
Single	24 (15)
Married	101 (63.1)
Divorced	28 (17.5)
Widowed	7 (4.4)
Religion	
Christian	122 (76.3)
Muslim	33 (20.6)
Traditionalist	2 (1.3)
None	3 (1.9)
Previous occupation	
Low income	133 (83.1)
Medium income	17 (10.6)
High income	10 (6.3)
Length of sentence	
<1 year	6 (3.8)
1-10 years	58 (36.2)
11-20 years	58 (36.2)
21-30 years	19 (11.9)
31-40 years	3 (1.9)
Life imprisonment	8 (5.0)
Remand	8 (5.0)
Length of stay	
<1 year	40 (25.0)
1-2 years	27 (16.9)
3-5 years	33 (20.6)
>5 years	60 (37.5)

*Data is presented in percentages and frequencies with percentages in parenthesis. *JHS means Junior High School and SHS means Senior Secondary School*

4. DISCUSSION

Male inmates dominated the study population and a majority of participants belonged to the lowest age of inclusion. This is consistent with the global data on the predominance of males on the penal system and the higher rates of crime among younger persons [17].

The prevalence of dyslipidaemia (57.5%) and hypertension (57.5%) were high among inmates and these can be attributed to inappropriate diet and high rates of physical inactivity observed [18,19]. Elevated serum LDL, reduced HDL and hypertension are prominent metabolic risk factors and predispose inmates to cardiovascular diseases [20]. Continual treatment and control of high blood pressure is essential in reducing

future cardiovascular risk [21] but only 11.9% of all those whose blood pressure satisfied the diagnostic criteria for hypertension had been diagnosed and out of this percentage 10.6% were on medication but none of them was on special diet. Meanwhile, the 2012 Ghana prisons service annual report indicated that

cardiovascular diseases are the leading cause of death among Ghanaian inmates. Other studies have also reported under diagnoses of diseases among prisoners [22,23] and this calls for regular health screening and improved health care for inmates.

Table 2. Means of metabolic characteristics of study participants

Parameters	Mean \pm SD	Gender		P-value
		Male	Female	
BMI (kg/m ²)	22.8 \pm 4.1	22.5 \pm 3.8	26.9 \pm 5.5	0.032*
Systolic(mm Hg)	141.1 \pm 23.2	140.5 \pm 23.7	149.3 \pm 11.4	0.050
Diastolic (mm Hg)	88.9 \pm 14	88.4 \pm 14.2	96.7 \pm 9.0	0.019*
FBG (mmol/L)	4.3 \pm 0.9	4.3 \pm 0.9	5.0 \pm 0.9	0.038*
HDL-c (mmol/L)	1.4 \pm 0.4	1.4 \pm 0.4	1.4 \pm 0.2	0.821
LDL-c (mmol/L)	3.0 \pm 1.0	2.9 \pm 1.0	3.4 \pm 0.8	0.151
Triglycerides (mmol/L)	1.1 \pm 0.6	1.1 \pm 0.7	0.9 \pm 0.3	0.045*
Total Cholesterol (mmol/L)	4.9 \pm 1.3	4.9 \pm 1.3	5.2 \pm 1.0	0.421
Waist Circumference (cm)	81.1 \pm 10.3	81.0 \pm 10.1	83.2 \pm 13.2	0.613

*Means of parameters are presented by gender. Independent sample T-test was used to compare males and females. *Significant differences exist at p<0.05*

Table 3. Prevalence of cardiovascular risk factors among inmates

Variables	Total n(%)	Male n(%)	Female n(%)	P-value
Blood Pressure				
Normotensive	22 (13.8)	22(14.7)	0(0)	0.094
Prehypertension	46 (28.8)	45 (30.0)	1(10)	
Hypertension	92 (57.5)	83 (55.3)	9 (90)	
Fasting blood glucose				
Hypoglycaemia	11 (6.9)	11 (7.3)	0 (0)	<0.001*
Normal blood glucose	142 (88.8)	135 (90.0)	7 (70)	
Pre-diabetes	5 (3.1)	2 (1.3)	3 (30)	
Diabetes	2 (1.2)	2 (1.3)	0 (0)	
BMI				
Underweight	11 (6.9)	11 (7.3)	0 (0)	0.005
Normal weight	113 (70.6)	109(72.7)	4 (40)	
Overweight	26 (16.2)	23 (15.3)	3 (30)	
Obese	10 (6.2)	7 (4.7)	3 (30)	
Waist circumference				
Normal waist circumference	149 (93.1)	143(95.3)	6 (60)	0.002*
Central obesity	11 (6.9)	7 (4.7)	4 (40)	
Lipid profile				
Normal lipid levels	67 (41.9)	63 (42.3)	4 (40)	1.000
Dyslipidaemia	92 (57.5)	86(57.7)	6 (60)	
Metabolic syndrome (> 2 parameters)	13 (8.1)	9 (6.0)	4 (40)	0.004*

Categorical data on cardiovascular risk factors. Gender groupings were compared using chi-square test.

**Significant differences exist at p<0.05*

Table 4. Physical activity and fruit intake by inmates

Variable	Frequency n(%)
Voluntary exercise	
Yes	72(45)
No	88(55)
Time spent on exercise	
<30 minutes	31(19.4)
30 minutes	20(12.5)
45 minutes	16(10)
>45 minutes	5(3.1)
Frequency of exercise	
Daily	29(8.1)
Weekly	19(11.9)
Monthly	2(1.3)
Occasionally	22(13.8)
Frequency of fruit intake	
Daily	3(1.6)
Weekly	10(6.3)
Monthly	9(5.6)
Occasionally	84(52.5)
Never	54(33.8)

Data is presented categorically with percentage in parenthesis

Table 5. Medical history of inmates

Variable	Frequency n(%)
Presence of chronic condition	
Yes	24(15)
No	136(85)
Type of chronic condition	
Hypertension	19(11.9)
Diabetes	3(1.9)
Dyslipidaemia	0(0)
Hypertension and diabetes	1(6)
Medication for chronic disease	
Yes	17(10.6)
No	7(4.4)
Special diet for chronic disease	
Yes	0(0)
No	24(15)

Data is presented categorically with percentage in parenthesis

Inmates who were overweight or obese by BMI constituted 22.6%. The prevalence of central obesity (6.9%) was low especially among male inmates. Most studies conducted in developed countries have found a higher mean BMI than what was found in this study. Leigey and Johnston [24] recorded a mean BMI of 28.8 among prisoners in the United States. Togas, Raikou [25] found a mean BMI of 25.68 among Greece prisoners.

The prevalence of diabetes was 1.2% and this may be linked to the low prevalence of obesity especially central obesity observed in this study. Other studies have however reported a relatively higher prevalence than what was found in this study. Bai, Befus [26] for instance reported a prevalence of 5.1%. Some inmates were hypoglycaemic (6.9%) and this can suggest improperly managed diabetes or hunger [27]. Hypoglycaemia is associated with aggressive behaviour and can interfere with the reformation process [28].

The proportion of inmates with metabolic syndrome was 8.1%. The prevalence of metabolic syndrome was higher among female inmates compared to males. Metabolic syndrome poses a double fold risk to the development of cardiovascular diseases and a five-fold risk to the development of diabetes mellitus [29]. The 8.1% prevalence of metabolic syndrome among the participants of this study is substantially lower than the 18% prevalence reported in a study among the free Ghanaian population by Akpalu [30]. Silverman-Retana, Lopez-Ridaura [31] however reported a 2.91% prevalence among inmates included in their study and this is less than what was found in this study. Fruits and vegetables are protective against metabolic risk conditions but inmates consumed these occasionally.

The mean BMI, mean diastolic and systolic blood pressure and mean FBG were significantly higher for female inmates. Female inmates present with more health challenges compared to their male counterparts. Prisons are male dominated environments with meal provision and routines that suit males better than females. The prison system should therefore be reformed for the benefit of females.

5. CONCLUSION

Inmates had a high prevalence of hypertension (57.5%) and dyslipidaemia (57.5%), but the prevalence of diabetes and obesity were low. Most inmates were sedentary and occasionally consumed fruits. Females had a higher prevalence of cardiovascular risk factors compared to their male counterparts. Frequent health screening, initiation of exercise programmes and appropriate dietary provision should commence within prisons as measures to curb metabolic risk factors among prisoners. Blood pressure check can be done routinely to detect cases of hypertension.

CONSENT

As per international standard or university standard, patient's written consent has been collected and preserved by the authors.

ETHICAL APPROVAL

As per international standard or university standard, written approval of Ethics committee has been collected and preserved by the authors.

ACKNOWLEDGEMENT

We are grateful to the Ghana Prisons Service for the assistance during data collection at the prison facilities.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. Global atlas on cardiovascular disease prevention and control: Policies, strategies, and interventions. Geneva: WHO; 2011. Available:www.who.int/cardiovascular_diseases/publications/atlas.../en/
2. Toth PP, Henriksson KM, Palmer MK. Metabolic syndrome and low-density lipoprotein cholesterol (LDL-C) goal attainment in the National Health and Nutrition Examination Survey (NHANES) (2003–2012); 2016.
3. Lorenzo C, Williams K, Hunt KJ, Haffner SM. The National Cholesterol Education Program–Adult Treatment Panel III, International Diabetes Federation, and World Health Organization definitions of the metabolic syndrome as predictors of incident cardiovascular disease and diabetes. *Diabetes Care*. 2007;30(1):8-13.
4. Binswanger IA, Krueger PM, Steiner JF. Prevalence of chronic medical conditions among jail and prison inmates in the United States compared with the general population. *Journal of Epidemiology and Community Health*. 2009;jech. 2009; 090662.
5. Hannan-Jones M, Capra S. Prevalence of diet-related risk factors for chronic disease in male prisoners in a high secure prison. *European Journal of Clinical Nutrition*. 2016;70(2):212-6.
6. Ünal V, Ünal EÖ, Çetinkaya Z, İmalı M, Gürler S, Koç S. Custody and prison deaths autopsied in Istanbul between 2010 and 2012. *Journal of Forensic and Legal Medicine*. 2016;39:16-21.
7. Binswanger IA, Krueger PM, Steiner JF. Prevalence of chronic medical conditions among jail and prison inmates in the United States compared with the general population. *Journal of Epidemiology & Community Health*. 2009;jech. 2009; 090662.
8. World Health Organization. Prisons and health. Regional Office of Europe, Copenhagen, Denmark; 2014.
9. Ahalt C, Trestman RL, Rich JD, Greifinger RB, Williams BA. Paying the price: The pressing need for quality, cost, and outcomes data to improve correctional health care for older prisoners. *Journal of the American Geriatrics Society*. 2013;61(11):2013-2019.
10. Berry JD, Dyer A, Cai X, Garside DB, Ning H, Thomas A, et al. Lifetime risks of cardiovascular disease. *New England Journal of Medicine*. 2012;366(4):321-9.
11. Michaud M, Balardy L, Moulis G, Gaudin C, Peyrot C, Vellas B, et al. Proinflammatory cytokines, aging, and age-related diseases. *Journal of the American Medical Directors Association*. 2013;14(12):877-82.
12. La Vigne N, Samuels J. The growth & increasing cost of the federal prison system: Drivers and potential solutions. Washington, DC: Urban Institute; 2012. Available:<http://www.urban.org/UploadedPDF/412693-The-Growth-and-Increasing-Cost-of-the-Federal-Prison-System.pdf>
13. Li SD. Toward a cost-effective correctional system: New developments in community-based corrections in China. *Victims & Offenders*. 2014;9(1):120-5.
14. Schmitt J, Warner K, Gupta S. The high budgetary cost of incarceration. Washington, DC: Center for Economic and Policy Research; 2010. Available:<http://www.cepr.net/documents/publications/incarceration-2010-06.pdf>
15. Expert Panel on Detection E. Executive summary of the Third Report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). *Jama*. 2001;285(19):2486.
16. Organization WH. BMI classification; 2015.

- Available:<http://www.who.int/bmi/index.jsp>. 2016
17. Enggist S, Møller L, Galea G, Udesen C. Prisons and health. World Health Organization; 2014.
 18. Cook EA, Lee YM, White BD, Gropper SS. The diet of inmates: An analysis of a 28-day cycle menu used in a large county jail in the state of Georgia. *Journal of Correctional Health Care*. 2015;21(4):390-9.
 19. Fawad DA, Saqib DM, Gul DAM, Jan DHU. Frequency of cardiovascular risk factors among prisoners. *Pakistan Heart Journal*. 2012;43(1-2).
 20. Acharjee S, Boden WE, Hartigan PM, Teo KK, Maron DJ, Sedlis SP, et al. Low levels of high-density lipoprotein cholesterol and increased risk of cardiovascular events in stable ischemic heart disease patients: A post-hoc analysis from the COURAGE Trial (Clinical outcomes utilizing revascularization and aggressive drug evaluation). *Journal of the American College of Cardiology*. 2013;62(20):1826-33.
 21. World Health Organization. A global brief on hypertension: Silent killer, global public health crisis. World; 2016.
 22. Vera-Remartínez E, Borraz-Fernández J, Domínguez-Zamorano J, Mora-Parra L, Casado-Hoces S, González-Gómez J, et al. Prevalence of chronic diseases and risk factors among the Spanish prison population. *Rev Esp Sanid Penit*. 2014;16(2):38-47.
 23. Enggist S, Møller L, Galea G, Udesen C. Prisons and health: WHO Regional Office for Europe; 2014.
 24. Leigey ME, Johnston ME. The prevalence of overweight and obesity among aging female inmates. *Journal of Correctional Health Care*. 2015;21(3):276-85.
 25. Togas C, Raikou M, Niakas D. An assessment of health related quality of life in a male prison population in Greece associations with health related characteristics and characteristics of detention. *BioMed Research International*; 2014.
 26. Bai JR, Befus M, Mukherjee DV, Lowy FD, Larson EL. Prevalence and predictors of chronic health conditions of inmates newly admitted to maximum security prisons. *Journal of Correctional Health Care*. 2015;21(3):255-64.
 27. Desouza CV, Bolli GB, Fonseca V. Hypoglycemia, diabetes, and cardiovascular events. *Diabetes care*. 2010;33(6):1389-94.
 28. Benton D. The impact of diet on anti-social, violent and criminal behaviour. *Neuroscience & Biobehavioral Reviews*. 2007;31(5):752-74.
 29. DeFina LF, Vega GL, Leonard D, Grundy SM. Fasting glucose, obesity, and metabolic syndrome as predictors of type 2 diabetes. *Journal of Investigative Medicine*. 2012;60(8):1164-8.
 30. Akpalu J, Akpalu A, Ofei F. The metabolic syndrome among patients with cardiovascular disease in Accra, Ghana. *Ghana Medical Journal*. 2011;45(4).
 31. Silverman-Retana O, Lopez-Ridaura R, Servan-Mori E, Bautista-Arredondo S, Bertozzi SM. Cross-sectional association between length of incarceration and selected risk factors for non-communicable chronic diseases in two male prisons of Mexico City. *PloS One*. 2015;10(9):e0138063.

© 2018 Agyapong et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/23830>*